

Dumfries & Galloway Royal infirmary  
RITUXIMAB for Rheumatoid Arthritis  
Rheumatology: Consultant Dr M J McMahon

Infusion Number  
**2 3 4 5 6**

Name  
Ref Number  
Ward 10

PRE-TREATMENT MEDICATIONS

			time	given by
PARACETAMOL	1000 mg PO	30 minutes before rituximab	_____	_____
CHLORPHENIRAMINE	10 mg IV	30 minutes before rituximab	_____	_____
METHYLPREDNISOLONE	100 mg IV	250 mls/hr, before rituximab	_____	_____
in 100 mls 0.9% Saline				

PRN Medications

CHLORPHENIRAMINE 10 mg IV tds  
PARACETAMOL 1000 mg PO 4 hourly  
HYDROCORTISONE 100 mg IV tds  
METOCLOPRMIDE 10 mg IV 8 hourly

Reduce rituximab infusion rate to 50% for minor infusion reactions.

RITUXIMAB 1000 mg in 500 mls 0.9% Saline

Batch no. _____	time	given by	checked
50 mls/hr for 30 minutes	_____	_____	_____
100 mls/hr for 30 minutes	_____	_____	_____
150 mls/hr for 30 minutes	_____	_____	_____
then 200 mls/hr	_____	_____	_____

PRESCRIBER

DATE

FOR USE ON

COMMENTS